

ADVIL- ibuprofen tablet, coated
GlaxoSmithKline Consumer Healthcare Holdings (US) LLC

Drug Facts

ACTIVE INGREDIENT

Advil Tablets (in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Advil Caplets (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Advil Gel Caplets (in each gel caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever/Fever reducer

USES

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis

- temporarily reduces fever

WARNINGS

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)

- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

Advil Tablets

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Advil Caplets

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Advil Gel Caplets

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 gel caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 gel caplet, 2 gel caplets may be used
- do not exceed 6 gel caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

OTHER INFORMATION

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

INACTIVE INGREDIENTS

Advil Tablets

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Advil Caplets

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Advil Gel Caplets

colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C red no. 40, FD&C yellow no. 6, fractionated coconut oil, gelatin, glycerin, hypromellose, pharmaceutical ink, pregelatinized starch, propyl gallate, purified water, sodium lauryl sulfate, stearic acid, synthetic iron oxides, titanium dioxide, triacetin

Questions or comments?

call toll free **1-800-88-ADVIL**

PRINCIPAL DISPLAY PANEL - 200 mg Gel Caplet Bottle Label

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

100 Gel Caplets

Gel Caplets

**READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION**

**Do Not Use if seal under bottle cap imprinted with
"SEALED for YOUR PROTECTION" is broken or missing.**

Advil[®]
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)
100 Gel Caplets **Gel Caplets**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses
 ■ temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis. ■ temporarily reduces fever

Warnings
 ■ Ask your doctor before use if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems.
 ■ This product may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Do not use this product if you have ever had an allergic reaction to any pain reliever/fever reducer.
 ■ Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions
 ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ adults and children 12 years and over: take 1 gel caplet every 4 to 6 hours while symptoms persist
 ■ if pain or fever does not respond to 1 gel caplet, 2 gel caplets may be used
 ■ do not exceed 6 gel caplets in 24 hours, unless directed by a doctor
 ■ children under 12 years: ask a doctor
 Questions or comments? call toll free: 1-800-68-ADVIL
 For most recent product information, visit www.Advil.com
 Pfizer, Madison, NJ 07940 USA
 © 2016 Pfizer Inc.

PAA080251

PRINCIPAL DISPLAY PANEL - 100 Count 200 mg Gel Caplet Bottle Carton


Advil[®]

**Ibuprofen Tablets, 200 mg
Pain Reliever / Fever Reducer (NSAID)**

100 Coated Gel Caplets*

**Capsule-Shaped Gelatin-Coated Tablets*

**Gel
Caplets**



PRINCIPAL DISPLAY PANEL - 50 Pouch Dispenser


Advil®

**Ibuprofen Tablets, 200 mg
Pain Reliever / Fever Reducer (NSAID)**

Tablets

1. **Headache**
2. **Minor Pain of Arthritis**
3. **Toothache**
4. **Muscular Aches**
5. **Menstrual Cramps**
6. **Fever**

50 PACKETS OF 2 COATED TABLETS EACH



PRINCIPAL DISPLAY PANEL - 300 Tablet Bottle Label - NDC 0573-0154-98

Advil®
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)
300 Coated Tablets

Advil®

**Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)**

**Advil®
Tablets**

300 Coated Tablets

PAA083046

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Drug Facts

Active ingredient (in each tablet)	Purpose
Ibuprofen 200 mg (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
 - temporarily reduces fever

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- coughing (wheezing)
- shock
- skin redness
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

NO COPY HINGED AREA

Drug Facts (continued)

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, polydioxane, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Questions or comments?
call toll free 1-800-89-ADVIL

Advil®

For most recent product information, visit www.Advil.com

Pfizer, Madison, NJ 07940 USA

© 2016 Pfizer Inc.

NO COPY HINGED AREA

NO COPY AREA

PRINCIPAL DISPLAY PANEL - 200 Count 200 mg Tablet Bottle Label

**New Easy Open
ARTHRITIS CAP**

**Arthritis Foundation
Ease of**

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (**NSAID**)
10 Coated Tablets

ADVILO® SAFETY SEALED ADVIL® SAFETY SEALED ADVILO® SAFETY SEALED

TEAR HERE TO OPEN

123 CODE FPO

Drug Facts (continued)

- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use Ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, polydioxane, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Questions or comments?

call toll free 1-800-88-ADWL

DO NOT USE IF SEAL AROUND CAP IS BROKEN OR MISSING. ADVIL SAFETY SEALED.

For most recent product information, visit www.Advil.com

Pfizer, Madison, NJ 07940 USA
© 2017 Pfizer Inc.

PAAD96199



Advil ADVIL Tablets

Ibuprofen Tablets, 200 mg
Pain Reliever /Fever Reducer (NSAID)
10 Coated Tablets

Drug Facts

Active ingredient (in each tablet)	Purpose
Ibuprofen 200 mg (NSAID)*	Pain reliever/ Fever reducer
*nonsteroidal anti-inflammatory drug	

LIFT HERE for More Drug Facts

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscle aches
 - minor pain of arthritis
 - temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ shock ■ skin redness ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

- The chance is higher if you
 - are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better





PRINCIPAL DISPLAY PANEL - 200 mg Tablet Vial Label

Advil[®]

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

10 Tablets

Tablets



PRINCIPAL DISPLAY PANEL - 10 Tablet Package

Pocket Pack

Do not use if blister is open or the words "ADVIL SAFETY SEALED" under blister are missing or torn.

Advil®

Ibuprofen Tablets, 200 mg

Pain Reliever / Fever Reducer (NSAID)

10 Coated Tablets

Tablets

Pocket Pack

Do not use if blister is open or the words "ADVIL SAFETY SEALED" under blister are missing or torn.

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

10 Coated Tablets

Tablets

READ AND KEEP CARD FOR COMPLETE WARNINGS AND INFORMATION

FPO
3 0573 0151 10 8

Drug Facts

Active ingredient (in each tablet): Purpose
Ibuprofen 200 mg (NSAID) Pain reliever/fever reducer
"nonsteroidal anti-inflammatory drug"

Uses

- temporarily relieves minor aches and pains due to:
 - headaches
 - backache
 - backache
 - muscular cramps
 - the common cold
 - menstrual aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness/itch
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Food use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

Drug Facts (continued)

When using this product:

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - belly hurt
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- read all warnings and directions before use. Keep cart.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients
acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmanolized ink, polydioxane, polydioxane starch, polypropylene glycol, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Questions or comments? call toll free 1-800-88-ADVIL

PAA063442

For most recent product information, visit www.Advil.com
Pfizer Inc., New York, NY 10040 USA © 2017 Pfizer Inc.

PRINCIPAL DISPLAY PANEL - 115 Tablet Bottle Label

Advil®

Ibuprofen Tablets, 200 mg

Pain Reliever/Fever Reducer (NSAID)

115 Tablets

Tablets

READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with
"SEALED for YOUR PROTECTION" is broken or missing.

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

115 Tablets Tablets

Uses

- temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis.
- temporarily reduces fever

Warnings

- **Ask your doctor before use** if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems.
- This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Do not use** this product if you have ever had an allergic reaction to any pain reliever/fever reducer.
- **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:
 - are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed.

Directions

- **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Questions or comments?
call toll free 1-800-88-ADVIL
For most recent product information, visit www.Advil.com
Pfizer, Madison, NJ 07940 USA
© 2016 Pfizer Inc.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

PAA083084




PRINCIPAL DISPLAY PANEL - 115 Tablet Bottle Carton - NDC 0573-0151-05

15
FREE
TABLETS

Advil®
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

115
Coated Tablets
Tablets



PRINCIPAL DISPLAY PANEL - 225 Caplet Bottle Label

Advil®
 Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer (NSAID)

225 Caplets

Caplets

READ AND KEEP CARTON FOR COMPLETE
 WARNINGS AND INFORMATION

Advil®
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

225 Caplets **Caplets**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses ■ temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis. ■ temporarily reduces fever

Warnings
 ■ Ask your doctor before use if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems.
 ■ This product may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. Do not use this product if you have ever had an allergic reaction to any pain reliever/fever reducer.
 ■ Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

■ Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

Questions or comments?
 call toll free 1-800-88-ADVIL
 For most recent product information, visit www.Advil.com
 Pfizer, Madison, NJ 07940 USA
 © 2017 Pfizer Inc.

PAA085439

PRINCIPAL DISPLAY PANEL - 225 Caplet Bottle Carton - NDC 0573-0161-85

25
 FREE
 CAPLETS


Advil®
 Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer (NSAID)

225
 Coated Caplets*
 *Capsule-Shaped Tablets
 Caplets




PRINCIPAL DISPLAY PANEL - 24 Tablet Bottle Label

Advil®
 Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer (NSAID)
 24 Tablets
 Tablets
 READ AND KEEP CARTON FOR COMPLETE
 WARNINGS AND INFORMATION



PRINCIPAL DISPLAY PANEL - 24 Tablet Bottle Carton - NDC 0573-0150-20

Advil[®]
 Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer (NSAID)
 24 Coated Tablets
 Tablets



PRINCIPAL DISPLAY PANEL - 100 Tablet Bottle Label

Advil®

**Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)**

**100 Tablets
Tablets**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.


PRINCIPAL DISPLAY PANEL - 100 Count 200 mg Tablet Bottle Carton

Advil®

**Ibuprofen Tablets, 200 mg
Pain Reliever / Fever Reducer (NSAID)**

100 Coated Tablets

Tablets



PRINCIPAL DISPLAY PANEL - 100 Tablet Bottle Label

Advil®
 Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer (NSAID)

100 Tablets

Tablets

READ AND KEEP CARTON FOR COMPLETE
 WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with
 "SEALED for YOUR PROTECTION" is broken or missing.

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

100 Tablets Tablets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses

- temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis. ■ temporarily reduces fever

Warnings

- **Ask your doctor before use** if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems.
- This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Do not use this product if you have ever had an allergic reaction to any pain reliever/fever reducer.**
- **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:
 - are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed.
- **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor


Questions or comments?
call toll free 1-800-88-ADVIL

For most recent product information, visit www.Advil.com
Pfizer, Madison, NJ 07940 USA
© 2016 Pfizer Inc.

PAA080142

PRINCIPAL DISPLAY PANEL - 100 Coated Tablet Bottle Carton - NDC 0573-0150-40

Advil®
 Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer (NSAID)
 100 Coated Tablets
 Tablets



PRINCIPAL DISPLAY PANEL - 24 Caplet Bottle Label

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

24 Caplets

Caplets

READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION



PRINCIPAL DISPLAY PANEL - 24 Caplet Bottle Carton - NDC 0573-0160-20

Advil[®]

Ibuprofen Tablets, 200 mg

Pain Reliever/Fever Reducer (NSAID)

24 Coated Caplets*

*Capsule-Shaped Tablets

Caplets

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each caplet) **Ibuprofen** 200 mg (NSAID)¹ Pain reliever/fever reducer
acetaminophen 325 mg Pain reliever/fever reducer

Uses

- temporarily relieve minor aches and pains due to:
 - headache
 - toothache
 - rheumatoid arthritis
 - menstrual cramps
 - the common cold
 - muscle aches
 - minor pain of arthritis
 - temporarily relieve fever

Warnings

Always read the label. Read every time you take. Save the label for reference. Especially if people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin redness
- rash
- blisters

 If an allergic reaction occurs, stop use and seek medical help right away.

PAA083047

Drug Facts (continued)

Some health warning: This product contains an NSAID, which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or celecoxib)
- have 3 or more alcohol drinks every day while using this product
- take more or for longer time than directed

 Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use:

- if you have ever had an allergic reaction to any of your pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver or kidney, kidney disease, asthma, or had a stroke
- you are taking a diuretic


Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs

For most recent product information, visit www.Advii.com
 Pfizer, Madison, NJ 07640 USA © 2016 Pfizer Inc.



Drug Facts (continued)

Stop use and ask doctor if:

- you experience any of the following signs of stomach bleeding:
 - red, black, or dark stools
 - stools that look like coffee grounds
 - you have stomach pain, indigestion or stomach ache
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless otherwise directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not take more than directed.

- The smallest effective dose should be used
- Adults and children 12 years and over: Take 1 caplet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 caplet, 2 caplets may be used. Do not exceed 6 caplets in 24 hours, unless directed by a doctor
- Children under 12 years: Ask a doctor



Advil[®]

Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer (NSAID)

24 Coated Caplets*
 *Capsule-Shaped Tablets



PRINCIPAL DISPLAY PANEL - 200 mg Caplet Bottle Label

Advil[®]

**Ibuprofen Tablets, 200 mg
 Pain Reliever/Fever Reducer (NSAID)**

**100 Caplets
 Caplets**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Advil®
Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)
100 Caplets **Caplets**

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" is broken or missing.

Uses ■ temporarily relieves minor aches and pains due to: headache, muscular aches, menstrual cramps, the common cold, backache, toothache, minor pain of arthritis.
■ temporarily reduces fever

Warnings
■ **Ask your doctor before use** if you are pregnant, under a doctor's care for a serious condition, age 60 or over, taking any other drug or have stomach problems.
■ This product may cause a **severe allergic reaction**, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Do not use** this product if you have ever had an allergic reaction to any pain reliever/fever reducer.
■ **Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed.

■ **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions
■ **do not take more than directed**
■ **the smallest effective dose should be used**
■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
■ children under 12 years: ask a doctor

Questions or comments?
call toll free 1-800-88-ADVIL
For most recent product information, visit www.Advil.com
Pfizer, Madison, NJ 07940 USA
© 2016 Pfizer Inc.

PAA080148

PRINCIPAL DISPLAY PANEL - 100 Count 200 mg Caplet Bottle Carton


Advil®

**Ibuprofen Tablets, 200 mg
Pain Reliever / Fever Reducer (NSAID)**

50 Coated Caplets*

***Capsule-Shaped Tablets**

Caplets



PRINCIPAL DISPLAY PANEL - 36 Caplet Bottle Label

Advil®

Ibuprofen Tablets, 200 mg
Pain Reliever/Fever Reducer (NSAID)

36 Caplets

Caplets

READ AND KEEP CARTON FOR COMPLETE
WARNINGS AND INFORMATION



PRINCIPAL DISPLAY PANEL - 36 Caplet Bottle Carton - NDC 0573-0160-25

50%

MORE

FREE

12 Free

Caplets

Advil®

Ibuprofen Tablets, 200 mg

Pain Reliever/Fever Reducer (NSAID)

36

Coated Caplets*

*Capsule-Shaped Tablets

Caplets



ADVIL			
ibuprofen tablet, coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0165
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	BROWN (one side is brown while the other side is yellow)	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	Advil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-0165-20	1 in 1 CARTON	05/18/1984	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0573-0165-30	1 in 1 CARTON	05/18/1984	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0573-0165-40	1 in 1 CARTON	05/18/1984	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0573-0165-41	1 in 1 CARTON	05/18/1984	
4		125 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0573-0165-42	1 in 1 CARTON	05/18/1984	
5		150 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:0573-0165-31	1 in 1 CARTON	05/18/1984	
6		75 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:0573-0165-25	1 in 1 CARTON	05/18/1984	

7	36 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	05/18/1984	

ADVIL

ibuprofen tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0154
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients	
Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
DIACETYLATED MONOGLYCERIDES (UNII: 5Z17386USF)	

Product Characteristics			
Color	BROWN (pinkish brown)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	Advil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-0154-35	1 in 1 CARTON	05/18/1984	
1		150 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0573-0154-75	1 in 1 CARTON	05/18/1984	
2		200 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0573-0154-17	3000 in 1 CASE	05/18/1984	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:0573-0154-89	50 in 1 CASE	05/18/1984	
4		2 in 1 POUCH; Type 0: Not a Combination Product		
5	NDC:0573-0154-84	1 in 1 CARTON	05/18/1984	
5		225 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:0573-0154-98	300 in 1 BOTTLE; Type 0: Not a Combination Product	05/18/1984	
7	NDC:0573-0154-16	3 in 1 CARTON	05/18/1984	
7		2 in 1 POUCH; Type 0: Not a Combination Product		
8	NDC:0573-0154-60	360 in 1 BOTTLE; Type 0: Not a Combination Product	05/18/1984	
9	NDC:0573-0154-59	1 in 1 CARTON	05/18/1984	
9		200 in 1 BOTTLE; Type 0: Not a Combination Product		
10	NDC:0573-0154-21	200 in 1 BOTTLE; Type 0: Not a Combination Product	01/04/2016	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	05/18/1984	

ADVIL

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0151
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	

PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
DIACETYLATED MONO GLYCERIDES (UNII: 5Z17386USF)	

Product Characteristics

Color	BROWN (pinkish brown)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	Advil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-0151-12	10 in 1 VIAL; Type 0: Not a Combination Product	05/18/1984	
2	NDC:0573-0151-13	3 in 1 PACKAGE	05/18/1984	
2		10 in 1 VIAL; Type 0: Not a Combination Product		
3	NDC:0573-0151-10	10 in 1 PACKAGE	05/18/1984	
3		10 in 1 VIAL; Type 0: Not a Combination Product		
4	NDC:0573-0151-21	1 in 1 CARTON	05/18/1984	
4		36 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0573-0151-20	2 in 1 PACKAGE	05/18/1984	
5		10 in 1 VIAL; Type 0: Not a Combination Product		
6	NDC:0573-0151-48	2 in 1 PACKAGE	05/18/1984	
6		1 in 1 CARTON		
6		100 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:0573-0151-30	1 in 1 CARTON	05/18/1984	
7		130 in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:0573-0151-05	1 in 1 CARTON	05/18/1984	
8		115 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	05/18/1984	

ADVIL

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0161
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
DIACETYLATED MONOGLYCERIDES (UNII: 5Z17386USF)	

Product Characteristics

Color	BROWN (pinkish brown)	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	Advil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-0161-35	1 in 1 CARTON	05/18/1984	
1		150 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0573-0161-51	1 in 1 CARTON	05/18/1984	
2		200 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0573-0161-85	1 in 1 CARTON	05/18/1984	
3		225 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0573-0161-65	65 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	05/18/1984	

ADVIL

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0166
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
WATER (UNII: 059QF0K00R)	

Product Characteristics

Color	BROWN (one side is brown while the other side is yellow)	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	Advil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:0573-0166-51	1 in 1 CARTON	05/18/1984	
1		200 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0573-0166-85	225 in 1 BOTTLE; Type 0: Not a Combination Product	05/18/1984	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	05/18/1984	

ADVIL

ibuprofen tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0150
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients	
Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
DIACETYLATED MONOGLYCERIDES (UNII: 5Z17386USF)	

Product Characteristics			
Color	BROWN (pinkish brown)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	Advil

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-0150-20	1 in 1 CARTON	05/18/1984	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0573-0150-30	1 in 1 CARTON	05/18/1984	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0573-0150-40	1 in 1 CARTON	05/18/1984	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0573-0150-31	1 in 1 CARTON	05/18/1984	
4		75 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0573-0150-32	1 in 1 CARTON	05/18/1984	
5		60 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:0573-0150-41	1 in 1 CARTON	05/18/1984	
6		125 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:0573-0150-42	1 in 1 CARTON	05/18/1984	
7		150 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	05/18/1984	

ADVIL

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-0160
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
METHYL PARABEN (UNII: A218C7H9T)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	

SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
DIACETYLATED MONO GLYCERIDES (UNII: 5Z17386USF)	

Product Characteristics

Color	BROWN (pinkish brown)	Score	no score
Shape	OVAL	Size	15mm
Flavor		Imprint Code	Advil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-0160-20	1 in 1 CARTON	05/18/1984	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0573-0160-30	1 in 1 CARTON	05/18/1984	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0573-0160-40	1 in 1 CARTON	05/18/1984	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:0573-0160-25	1 in 1 CARTON	05/18/1984	
4		36 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:0573-0160-31	1 in 1 CARTON	05/18/1984	
5		75 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:0573-0160-41	1 in 1 CARTON	05/18/1984	
6		125 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:0573-0160-42	1 in 1 CARTON	05/18/1984	
7		150 in 1 BOTTLE; Type 0: Not a Combination Product		
8	NDC:0573-0160-17	1 in 1 CARTON	05/18/1984	
8		115 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	05/18/1984	

ADVIL
 ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0573-1711
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
WHITE WAX (UNII: 7G1J5DA97F)	
DIACETYLATED MONOGLYCERIDES (UNII: 5Z17386USF)	

Product Characteristics

Color	BROWN (pinkish brown)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	Advil
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0573-1711-14	1 in 1 CARTON	05/18/1984	
1		225 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	05/18/1984	

Labeler - GlaxoSmithKline Consumer Healthcare Holdings (US) LLC (079944263)**Establishment**

Name	Address	ID/FEI	Business Operations
Wyeth Pharmaceuticals Company		829390975	ANALYSIS(0573-0150, 0573-0151, 0573-0154, 0573-0160, 0573-0161, 0573-0165, 0573-0166, 0573-1711), LABEL(0573-0150, 0573-0151, 0573-0154, 0573-0160, 0573-0161, 0573-0165, 0573-0166, 0573-1711), MANUFACTURE(0573-0150, 0573-0151, 0573-0154, 0573-0160, 0573-0161, 0573-0165, 0573-0166, 0573-1711), PACK(0573-0150, 0573-0151, 0573-0154, 0573-0160, 0573-0161, 0573-0165, 0573-0166, 0573-1711)

Revised: 11/2020

GlaxoSmithKline Consumer Healthcare Holdings (US) LLC